

SUPPLEMENTAL TABLE 1
Polymerase chain reaction (PCR) primers used in this study

Target locus	Primer/probe	Sequence	Refs.
SYBR-based nested PCR			
<i>Cryptosporidium</i> SSU rRNA (primary amplification)	Forward	TTCTAGAGCTAATACATGCG	20
	Reverse	CCCATTTCCTTCGAAACAGGA	
<i>Cryptosporidium</i> SSU rRNA (secondary amplification)	Forward	GGAAGGGTTGTATTTATTAGATAAAG	
	Reverse	AAGGAGTAAGGAACAACCTCCA	
Probe-based assay			
<i>Cryptosporidium parvum</i> SSU rRNA	Forward	TCCTTGAAATGAATATTTGTGACTCG	21
	Reverse	TTAATGTGGTAGTTGCGGTTGAAC	
<i>Cryptosporidium hominis</i> SSU rRNA	Probe	FAM-TATCTCTTCGTAGCGGCGTA-BHQ1	21
	Forward	TCCTTGAAATGAATATTTGTGACTCG	
	Reverse	AAATGTGGTAGTTGCGGTTGAAA	
<i>Cryptosporidium meleagridis</i> 60 kDa glycoprotein 60	Probe	VIC-CTTACTTCGTGGCGGCGT-BHQ1	Present study
	Forward	GAGCTCAGCACTCTCTACTA	
	Reverse	GCGTCTGTGAGTGATCTTCTT	
	Probe	TEX615-ATGACACTGCGTCTGAGGAAAGCT-3IAbrQSp	

SSU = small subunit.